

BOUNDARY VALUE PROBLEMS FOR  $\nabla^2 u + k^2 u = 0$

BY

IVAR STAKGOLD

TECHNICAL REPORT NO. 17

JANUARY 29, 1954

PREPARED UNDER CONTRACT Nonr-225(11)

(NR-041-036)

FOR

OFFICE OF NAVAL RESEARCH

APPLIED MATHEMATICS AND STATISTICS LABORATORY

STANFORD UNIVERSITY

STANFORD, CALIFORNIA

# A BOUNDARY VALUE PROBLEM FOR $\nabla^2 u + k^2 u = 0$

By

Ivar Stakgold

Stanford University<sup>1/</sup>

## Introduction and summary.

Part I: The solution of  $u_{xx} + u_{yy} + k^2 u = 0$ , with  $u$  (or  $\frac{\partial u}{\partial r}$ ) on a half-line is obtained by a simpler and more direct method than those previously found in the literature. The method applies equally well to a wedge of arbitrary opening. An idea of Magnus (1) is used to simplify the solution when the boundary values can be expanded in a series of Bessel functions.

Part II: The problem of diffraction by an infinite cone with a spherical tip is treated by a method similar to the used in part I. Both point source and plane wave excitations are considered. When the cone opening  $\theta_0$  is  $\frac{\pi}{2}$ , the problem reduces to reflection from a plane with a hemispherical boss. This simpler problem has been solved by Twersky (7); our results agree with his solution.

## Part I.

(a) We first determine the Green's function for the problem.

A two-dimensional unit source is located at the point  $(\rho', \theta')$  in front of a semi-infinite sheet ( $\theta = 0$ ).

We wish to solve the two-dimensional problem

---

<sup>1/</sup> On leave from Harvard University.

$$(1) \quad \frac{1}{\rho} \frac{\partial}{\partial \rho} \left( \rho \frac{\partial u}{\partial \rho} \right) + \frac{1}{\rho^2} \frac{\partial^2 u}{\partial \theta^2} + k^2 u = - \frac{\delta(\rho - \rho') \delta(\theta - \theta')}{\rho'}$$

$$u(\rho, 0+) = 0$$

$$u(\rho, 2\pi-) = 0 \quad .$$

$u$  obeys the Sommerfeld radiation condition at infinity

$$\lim_{r \rightarrow \infty} \sqrt{r} \left( \frac{\partial u}{\partial r} - iku \right) = 0$$

( $r$  is measured from an arbitrary fixed origin).

It is immediately clear that the angular eigenfunctions of the homogeneous equation are just  $\sin \frac{n}{2} \theta$ . This is a complete orthogonal set of functions over  $0 < \theta < 2\pi$ , satisfying the boundary conditions.

We write  $u(\rho, \theta) = \sum_{n=1}^{\infty} a_n(\rho) \sin \frac{n}{2} \theta$  where the  $a_n(\rho)$  are to be determined.

We multiply equation (1) by  $\sin \frac{n}{2} \theta$  and integrate from 0 to  $2\pi$ , obtaining:

$$\begin{aligned} \pi a_n''(\rho) + \frac{\pi}{\rho} a_n'(\rho) + k^2 a_n(\rho) + \frac{1}{\rho} \int_0^{2\pi} \sin \frac{n\theta}{2} \frac{\partial^2 u}{\partial \theta^2} d\theta \\ = \frac{-\sin \frac{n}{2} \theta'}{\rho'} \delta(\rho - \rho') \end{aligned}$$

(A priori it is possible that the coefficients  $a_n(\rho)$  are no better than  $O(\frac{1}{n^2})$  for large  $n$ , therefore we may not be able to differentiate twice termwise with respect to  $\theta$ ).

Integrating twice by parts and using the boundary conditions on  $u$  and  $\sin \frac{n\theta}{2}$ , we readily obtain

$$a_n''(\rho) + \frac{a_n'(\rho)}{\rho} + (k^2 - \frac{n^2}{4}) a_n(\rho) = \frac{-\sin \frac{n}{2} \theta'}{\pi \rho'} \delta(\rho - \rho') .$$

Hence

$$a_n(\rho) = \begin{cases} A H_{\frac{n}{2}}^{(1)}(k\rho) & \rho > \rho' \\ B J_{\frac{n}{2}}(k\rho) & \rho < \rho' \end{cases} .$$

This satisfies the radiation condition at infinity and the finiteness condition at the origin.

We now require

$$a_n(\rho' +) = a_n(\rho' -), \quad \frac{da_n}{d\rho}(\rho' +) - \frac{da_n}{d\rho}(\rho' -) = \frac{-\sin \frac{n}{2} \theta'}{\pi \rho'} .$$

Recalling the Wronskian relation:

$$\rho' [H_{\frac{n}{2}}^{(1)}(k\rho') J_{\frac{n}{2}}(k\rho') - H_{\frac{n}{2}}^{(1)}(k\rho') J_{\frac{n}{2}}'(k\rho')] = \frac{2i}{\pi k} .$$

We have:

$$a_n(\rho) = \frac{i}{2} \sin \frac{n}{2} \theta' \begin{cases} H_{\frac{n}{2}}^{(1)}(k\rho) J_{\frac{n}{2}}(k\rho') & \rho > \rho' \\ H_{\frac{n}{2}}^{(1)}(k\rho') J_{\frac{n}{2}}(k\rho) & \rho < \rho' \end{cases}$$

We use the standard abbreviation:

$$a_n(\rho) = \frac{i}{2} \sin \frac{n}{2} \theta' H_{\frac{n}{2}}^{(1)}(k\rho >) J_{\frac{n}{2}}(k\rho <). \quad \text{The symbol } \rho > \text{ means } \begin{cases} \rho & \text{when } \rho > \rho' \\ \rho' & \text{when } \rho < \rho' \end{cases}.$$

Therefore we have the Green's function:

$$(2) \quad u(\rho, \theta) = G(\rho, \theta; \rho', \theta') = \frac{i}{2} \sum_{n=1}^{\infty} \sin \frac{n}{2} \theta \sin \frac{n}{2} \theta' H_{\frac{n}{2}}^{(1)}(k\rho >) J_{\frac{n}{2}}(k\rho <)$$

(Formulas of this general nature, derived by the use of the idea of sources on a Riemann surface, are to be found in A. Sommerfeld (2)).

Expression (2) may be rewritten

$$\begin{aligned} G(\rho, \theta; \rho', \theta') &= \frac{i}{2} \sum_{n=1}^{\infty} \sin n\theta \sin n\theta' H_n^{(1)}(k\rho >) J_n(k\rho <) \\ &+ \frac{i}{2} \sum_{n=0}^{\infty} \sin \frac{2n+1}{2} \theta \sin \frac{2n+1}{2} \theta' H_{\frac{2n+1}{2}}^{(1)}(k\rho >) J_{\frac{2n+1}{2}}(k\rho <) . \end{aligned}$$

It is easy to recognize the first series. Had we carried out a similar analysis for a completely infinite sheet, the Green's function would turn out to be  $i \sum_{n=1}^{\infty} \sin n\theta \sin n\theta' H_n^{(1)}(k\rho >) J_n(k\rho <)$ .

We have therefore

$$\frac{i}{2} \sum_{n=1}^{\infty} \sin n\theta \sin n\theta' H_n^{(1)}(k\rho >) J_n(k\rho <) = \frac{i}{8} \left\{ H_0^{(1)}(kR) - H_0^{(1)}(kR^*) \right\}$$

where  $R$  is the distance between the source and the observation point,

$R^*$  is the distance between the reflected source about  $\theta = 0$  and the observation point. This result could have been obtained from the familiar addition theorem for cylinder functions. We may write

$$(3) \quad G(\rho, \theta; \rho', \theta') = \frac{1}{8} \left\{ H_0^{(1)}(kR) - H_0^{(1)}(kR^*) \right\} \\ + \frac{i}{2} \sum_{n=0}^{\infty} \sin \frac{2n+1}{2} \theta \sin \frac{2n+1}{2} \theta' H_{\frac{2n+1}{2}}^{(1)}(k\rho >) J_{\frac{2n+1}{2}}(k\rho <) .$$

We are now in a position to solve the problem of diffraction from a half-plane. Let the incoming field be  $e^{ikx}$  propagating towards increasing  $x$ .




Fig. 1

The half-plane is located in the plane  $x = 0$ , with  $y \geq 0$ . We use the coordinate system indicated in Fig. 1. The total field is to vanish for  $x = 0, y \geq 0$ .

We can think of  $e^{ikx}$  as being the limiting case of a point source removed to  $x = -\infty$ .

Indeed suppose there is a two-dimensional unit source at  $\theta' = \frac{\pi}{2}$ ,  $\rho = \rho'$ .




Fig. 2

The field of this point source is just  $\frac{i}{4} H_0^{(1)}(kR)$ .

We have:

$$R = \frac{(\rho' - \rho \sin \theta)}{\cos \theta} = \frac{\rho' + x}{\cos \theta} .$$

When  $\rho'$  is taken very large, the asymptotic expansion for  $\frac{i}{4} H_0^{(1)}(kR)$  yields

$$\frac{i}{4} \sqrt{\frac{2}{\pi kR}} e^{-\frac{i\pi}{4}} e^{ikR} .$$

To obtain the limit  $e^{ikx}$  as  $\rho' \rightarrow \infty$ , consider

$$\frac{4}{i} \sqrt{\frac{\pi k \rho'}{2}} e^{\frac{i\pi}{4}} e^{-ik\rho'} \left\{ \frac{i}{4} H_0^{(1)}(kR) \right\} .$$

As  $\rho' \rightarrow \infty$ ,  $\cos \beta \rightarrow 1$ ,  $R - \rho' \rightarrow 0$ , and it is easy to see that the limit of the above expression is  $e^{ikx}$ .

It is reasonable to suppose that performing the same limiting process on (3) will yield the diffraction a plane wave normally incident on a half plane. The result is derived rigorously in (b).

Thus, the total field is given by:

$$u(\rho, \theta) = \lim_{\rho' \rightarrow \infty} \frac{4}{i} \sqrt{\frac{\pi k \rho'}{2}} e^{\frac{i\pi}{4}} e^{-ik\rho'} \left[ \frac{i}{8} \left\{ H_0^{(1)}(kR) - H_0^{(1)}(kR^*) \right\} \right. \\ \left. + \frac{i}{2} \sum_{n=0}^{\infty} \sin \frac{2n+1}{2} \theta \sin \frac{2n+1}{2} \frac{\pi}{2} H_{\frac{2n+1}{2}}^{(1)}(k\rho') J_{\frac{2n+1}{2}}(k\rho') \right] .$$

The terms outside the summation yield  $\frac{e^{ikx} - e^{-ikx}}{2}$ .

In the summation,  $\rho < \rho'$  and the asymptotic expansion for  $H_{\frac{2n+1}{2}}^{(1)}(k\rho')$  is used:

$$H_{\frac{2n+1}{2}}^{(1)}(k\rho') \sim \sqrt{\frac{2}{\pi k \rho'}} e^{ik\rho'} e^{-in\pi} .$$

We obtain

$$(4) \quad u(\rho, \theta) = i \sin kx + 2e^{-\frac{i\pi}{4}} \sum_{n=0}^{\infty} (i)^{-n} \sin \frac{2n+1}{2} \theta \sin \frac{2n+1}{2} \frac{\pi}{2} J_{\frac{2n+1}{2}}(k\rho) .$$

This formula can easily be identified with that of Magnus (1) who uses the time dependence  $e^{i\omega t}$  and a rotated coordinate system.

The results for the case where the normal derivative vanishes on the half-line are easily found:

$$(3a) \quad G(\rho, \theta; \rho', \theta') = \frac{i}{8} \left\{ H_0^{(1)}(kR) + H_0^{(1)}(kR^*) \right\} \\ + \frac{1}{2} \sum_{n=0}^{\infty} \cos \frac{2n+1}{2} \theta \cos \frac{2n+1}{2} \theta' H_{\frac{2n+1}{2}}^{(1)}(k\rho >) J_{\frac{2n+1}{2}}(k\rho <)$$

$$(4a) \quad u(\rho, \theta) = \cos kx + 2e^{-\frac{i\pi}{4}} \sum_{n=0}^{\infty} (i)^{-n} \cos \frac{2n+1}{2} \theta \cos \frac{2n+1}{2} \frac{\pi}{2} J_{\frac{2n+1}{2}}(k\rho) .$$

It is clear that the case of oblique incidence is easily handled by the same method.

(b) Consider the following two-dimensional boundary value problem:

$$\nabla^2 u + k^2 u = 0 \quad u(\rho, \theta+) = u(\rho, 2\pi-) = \text{given function } f(\rho);$$

$u$  is continuous throughout the plane;  $\text{grad } u$  is continuous everywhere except possibly on the half-line  $\theta = 0$ ;  $u$  obeys a suitable condition at infinity.



---

Remark on the condition at infinity.

The condition at infinity is not the Sommerfeld radiation condition

$$\lim_{\rho \rightarrow \infty} \sqrt{\rho} \left( \frac{\partial u}{\partial \rho} - iku \right) = 0 .$$

In the case of diffraction by an obstacle having some infinite dimension, neither the total field nor the scattered field can be expected to satisfy the radiation condition.

As a simple example, consider total reflection of a plane wave by an infinite sheet.

The incident field is  $e^{ikx}$ .

$$\text{The total field is } \begin{cases} e^{ikx} - e^{-ikx} & x < 0 \\ 0 & x > 0 \end{cases} .$$

The "scattered" field is  $-e^{ik|x|}$ . None of these fields obeys the Sommerfeld condition. The same is true for diffraction by a semi-infinite sheet.

This is not fatal, however, since all we really need is

$$(5) \quad \lim_{\rho \rightarrow \infty} \int_S \left\{ H_0^{(1)}(kR) \frac{\partial u}{\partial n} - u \frac{\partial}{\partial n} H_0^{(1)}(kR) \right\} d\sigma = 0 .$$

The "scattered" field in the problem of total reflection obeys condition (5).

In general, the geometrical optics field (i.e., no diffraction) will satisfy (5) (see D. S. Jones (3)). Since the scattered field is expected to differ from the geometrical optics field only by a field satisfying the Sommerfeld condition, (5) will hold for the scattered field of the half-plane problem. In any event we adopt (5) as the imposed condition at infinity. It is clear that if (5) holds with the free space Green's function  $H_0^{(1)}$ , it would also hold with the Green's function  $G$  of part I.

---

Let us apply Green's theorem to  $u$  and the Green's function  $G$  given by (2). As the boundary of the region of integration is removed to infinity, we obtain (in virtue of (5)):

$$\begin{aligned} u(\rho', \theta') &= \int_0^\infty d\rho \frac{f(\rho)}{\rho} \left\{ \frac{\partial u}{\partial \theta} \Big|_{\theta=0+} - \frac{\partial u}{\partial \theta} \Big|_{\theta=2\pi-} \right\} \\ &= \int_0^\infty \frac{d\rho}{\rho} f(\rho) \left\{ \sum_{n=0}^\infty \frac{1}{2} (2n+1) \sin \frac{2n+1}{2} \theta' H_{\frac{2n+1}{2}}^{(1)}(k\rho >) \right\} . \end{aligned}$$

Under suitable conditions, this may be written as

$$(6) \quad u(\rho', \theta') = \frac{1}{2} \sum_{n=0}^\infty (2n+1) \sin \frac{2n+1}{2} \theta' \int_0^\infty d\rho \frac{f(\rho)}{\rho} H_{\frac{2n+1}{2}}^{(1)}(k\rho >) J_{\frac{2n+1}{2}}(k\rho <) .$$

At first, it might appear that  $\lim_{\theta' \rightarrow 0+} u(\rho', \theta') = 0$ ; this is not true.

We will show that the integral in (6) is no better than  $O(\frac{1}{n})$ , hence the coefficient of  $\sin \frac{2n+1}{2} \theta'$  is no better than  $O(\frac{1}{n})$ ; the series will exhibit discontinuities at  $\theta' = 0, 2\pi$ , and  $\lim_{\theta' \rightarrow \begin{cases} 0+ \\ 2\pi- \end{cases}} u(\rho', \theta') = f(\rho')$ .

Formula (6) gives the solution of our problem in a rather untractable form. Magnus (1) noted that integrals of the type (6) are considerably simplified when  $f(\rho)$  admits a series development in Bessel functions of integral order. Let

$$I_{m,n} = \int_0^\infty \frac{J_m(k\rho)}{\rho} H_{\frac{2n+1}{2}}^{(1)}(k\rho >) J_{\frac{2n+1}{2}}(k\rho <) d\rho .$$

We split the integral into integrals from 0 to  $\rho'$  and  $\rho'$  to  $\infty$ .

In each integral we let  $t = k\rho$

$$I_{m,n} = \frac{t \left\{ J_m(t) \frac{J_{2n+1}(t)}{2} - \frac{J_{2n+1}(t)}{2} J_m(t) \right\}}{m^2 - \left(\frac{2n+1}{2}\right)^2} \Bigg|_0^{k\rho'} H_{\frac{2n+1}{2}}^{(1)}(k\rho') + \frac{t \left\{ J_m(t) \frac{H_{2n+1}^{(1)}(t)}{2} - \frac{H_{2n+1}^{(1)}(t)}{2} J_m(t) \right\}}{m^2 - \left(\frac{2n+1}{2}\right)^2} \Bigg|_{k\rho'}^{\infty} J_{\frac{2n+1}{2}}(k\rho') .$$

The contribution from 0 vanishes, the contribution at infinity is

$$\frac{2i}{\pi} e^{\frac{i\pi}{2} \left(m - \frac{2n+1}{2}\right)} \frac{1}{m^2 - \left(\frac{2n+1}{2}\right)^2} .$$

If we use in addition the formula for the Wronskian of part I, we obtain

$$I_{m,n} = -\frac{2i}{\pi} e^{-\frac{i\pi}{4}} \frac{(i)^{m-n}}{m^2 - \left(\frac{2n+1}{2}\right)^2} J_{\frac{2n+1}{2}}(k\rho') + \frac{2i}{\pi} \frac{J_m(k\rho')}{m^2 - \left(\frac{2n+1}{2}\right)^2} .$$

For the case  $f(\rho) = J_m(\rho)$  (6) reduces to:

$$u_m(\rho', \theta') = \frac{(i)^m e^{-\frac{i\pi}{4}}}{\pi} \sum_{n=0}^{\infty} (i)^{-n} \frac{(2n+1) \sin \frac{2n+1}{2} \theta'}{m^2 - \left(\frac{2n+1}{2}\right)^2} J_{\frac{2n+1}{2}}(k\rho') - \frac{J_m(k\rho')}{\pi} \sum_{n=0}^{\infty} \frac{2n+1}{m^2 - \left(\frac{2n+1}{2}\right)^2} \sin \frac{2n+1}{2} \theta' .$$

The first of these Fourier series in  $\theta'$  has coefficients  $O(\frac{1}{n^2})$  due to the rapid decrease with  $n$  of  $J_{\frac{2n+1}{2}}(k\rho')$ . This series represents a continuous function which vanishes at  $\theta' = 0, 2\pi$ . Hence the limiting values as  $\theta' \rightarrow 0+, 2\pi -$  are also zero.

The second series has coefficients  $O(\frac{1}{n})$  and may exhibit a discontinuity at  $\theta' = 0$ . As a matter of fact it can be shown by an elementary computation that

$$-\pi \cos m\theta = \sum_{n=0}^{\infty} \frac{2n+1}{m^2 - (\frac{2n+1}{2})^2} \sin \frac{2n+1}{2} \theta' .$$

Thus:

$$\lim_{\theta' \rightarrow \begin{cases} 0+ \\ 2\pi - \end{cases}} u_m(\rho', \theta') = J_m(k\rho')$$

as required. We can write:

$$u_m(\rho', \theta') = \frac{(i)^m e^{-\frac{i\pi}{4}}}{\pi} \sum_{n=0}^{\infty} (i)^{-n} \frac{(2n+1) \sin \frac{2n+1}{2} \theta'}{m^2 - (\frac{2n+1}{2})^2} J_{\frac{2n+1}{2}}(k\rho') + J_m(k\rho') \cos m\theta'.$$

In general if  $f(\rho)$  admits an expansion  $f(\rho) = \sum_{m=0}^{\infty} \beta_m J_m(k\rho)$  we will obtain

$$(7) \quad u(\rho', \theta') = \sum_{m=0}^{\infty} \beta_m J_m(k\rho') \cos m\theta' + \frac{e^{-\frac{i\pi}{4}}}{\pi} \sum_{m,n=0}^{\infty} \beta_m (i)^{m-n} (2n+1) \frac{\sin \frac{2n+1}{2} \theta'}{m^2 - (\frac{2n+1}{2})^2} J_{\frac{2n+1}{2}}(k\rho') .$$

It is now easy to solve the problem of diffraction by a half-plane. For simplicity we take normal incidence. The total field  $W(\rho, \theta)$  may be split up in an incident field  $e^{ikx}$  and a scattered field  $u(\rho, \theta)$  satisfying (5). The values of  $u(\rho, \theta)$  on the half-plane are just  $f(\rho) = -1$ . Since  $-1 = -\sum_{p=1}^{\infty} \epsilon_p J_{2p}(k\rho)$ , we can use (7), with  $\beta_m = -\epsilon_m (-i)^m \cos m \frac{\pi}{2}$  ( $\epsilon_0 = 1$ ,  $\epsilon_m = 2$ ,  $m > 1$ ), thus obtaining

$$u(\rho', \theta') = \sum_{m=0}^{\infty} \epsilon_m J_{2m}(k\rho') \cos 2m\theta' + \sum_{n=0}^{\infty} \frac{e^{-\frac{i\pi}{4}}}{\pi} (i)^{-n} (2n+1) \sin \frac{2n+1}{2} \theta' J_{\frac{2n+1}{2}}(k\rho') \sum_{m=0}^{\infty} -\epsilon_m \frac{\cos m \frac{\pi}{2}}{m^2 - (\frac{2n+1}{2})^2}.$$

The first sum is recognized as  $-\cos(k\rho' \sin \theta') = -\cos kx'$ . The last sum can be evaluated by an elementary computation (see Magnus (1), p. 175)

$$\sum_{m=0}^{\infty} -\epsilon_m \frac{\cos m \frac{\pi}{2}}{m^2 - (\frac{2n+1}{2})^2} = \frac{2}{2n+1} \sin \left( \frac{2n+1}{2} \frac{\pi}{2} \right).$$

Thus we have a single summation for  $u(\rho', \theta')$  and we obtain


$$(8) \quad W(\rho, \theta) = i \sin kx + 2e^{-i \frac{\pi}{4}} \sum_{n=0}^{\infty} (i)^{-n} \sin \frac{2n+1}{2} \theta \sin \frac{2n+1}{2} \frac{\pi}{2} J_{\frac{2n+1}{2}}(k\rho)$$

which agrees with formula (4) of (a).

It has been shown that (8) can be transformed into the Sommerfeld solution (see Magnus (1), p. 179) and into the Kontorowich-Lebedev solution ((4), p. 241).

## Part II.

We consider now the diffraction by a cone with a spherical tip.



The problem of the cone without a tip has been treated by Sollfrey (5).

Fig. 3

The cone is infinite and the source is taken at  $\theta = \pi$ ,  $r = r'$ . The problem can be solved with only slight additional difficulty for other locations of the source.

The total field satisfies

$$\nabla^2 u + k^2 u = \frac{-\delta(r - r') \delta(\theta)}{2\pi r^2 \sin \theta} \quad (\text{the field is independent of } \varphi)$$

The boundary condition is taken as  $u = 0$  on both the sphere and cone.

$$(9) \quad \frac{1}{r^2} \frac{\partial}{\partial r} (r^2 \frac{\partial u}{\partial r}) + \frac{1}{r^2 \sin \theta} \frac{\partial}{\partial \theta} (\sin \theta \frac{\partial u}{\partial \theta}) + k^2 u = \frac{-\delta(r - r') \delta(\theta)}{2\pi r^2 \sin \theta}.$$

When  $r \neq r'$ , the homogeneous equation can readily be separated yielding the equation for the  $\theta$  part of the separable solutions

$$\frac{1}{\sin \theta} \frac{\partial}{\partial \theta} (\sin \theta w') + \lambda w = 0 \quad \begin{array}{l} w(\theta_0) = 0 \\ w(\pi) \text{ finite} \end{array}$$

$\lambda$  is a separation constant. If we let  $\lambda = \nu(\nu+1)$ ,  $x = \cos \theta$ ,

$$\frac{d}{dx} [(1-x^2) \frac{dw}{dx}] + \nu(\nu+1)w = 0 \quad \begin{array}{l} w(-1) = \text{finite} \\ w(\cos \theta_0) = 0 \end{array}$$

The only solution which is finite at -1 is  $P_\nu(-x)$ , the Legendre function of the first kind.

The eigenvalues  $\nu$  are determined from the equation  $P_\nu(-\cos \theta_0) = 0$ . The functions  $P_\nu(-\cos \theta)$  form a complete orthogonal set in the interval  $\theta_0 < \theta < \pi$ , with weight  $\sin \theta$ . The normalization factor

$$\int_{\theta_0}^{\pi} P_\nu^2(-\cos \theta) \sin \theta d\theta = A_\nu$$

will be evaluated later.

We return to equation (9) and consider for  $u$  an expansion of the form

$$u = \sum_{\nu} a_{\nu}(r) P_{\nu}(-\cos \theta)$$

We multiply (9) by  $r^2 P_{\mu}(-\cos \theta) \sin \theta$  and integrate from  $\theta_0$  to  $\pi$ .

We obtain the differential equation for  $a_{\mu}(r)$ :

$$(10) \quad \frac{d}{dr} (r^2 a'_{\mu}) + k^2 r^2 a_{\mu} - \mu(\mu+1) a_{\mu} = \frac{-\delta(r-r')}{2\pi A_{\nu}}$$

The solution of (10) may be written as:

$$a_{\mu}(r) = C \frac{H_{\mu+1/2}(kr>)}{\sqrt{r>}} \frac{Z_{\mu+1/2}(kr<)}{\sqrt{r<}}$$

where  $Z_{\mu+1/2}$  is a cylinder function which vanishes at  $r = a$ . We have taken care of the boundary conditions at  $r = a$ ,  $r = \infty$ , and of continuity at  $r = r'$ . We determined  $C$  from the jump condition

$$a'_{\mu}(r'+) - a'_{\mu}(r'-) = - \frac{1}{2\pi r'^2}$$

$$a'_{\mu}(r'+) = C \frac{Z_{\mu+1/2}(kr')}{\sqrt{r'}} \cdot \left\{ \frac{k}{\sqrt{r'}} H_{\mu+1/2}^{(1)}(kr') - H_{\mu+1/2}^{(1)}(kr') \frac{1}{2r'^{3/2}} \right\}$$

$$a'_{\mu}(r'-) = C \frac{H_{\mu+1/2}^{(1)}(kr')}{\sqrt{r'}} \cdot \left\{ \frac{k}{\sqrt{r'}} Z_{\mu+1/2}(kr') - Z_{\mu+1/2}(kr') \frac{1}{2r'^{3/2}} \right\} .$$

Hence

$$C \frac{k}{r'} \text{Wronskian} \left\{ Z_{\mu+1/2}, H_{\mu+1/2}^{(1)} \right\} = - \frac{1}{2\pi r'^2 A_{\nu}} .$$

We write

$$Z_{\mu+1/2}(kr) = H_{\mu+1/2}^{(1)}(kr) - \frac{H_{\mu+1/2}^{(1)}(ka)}{H_{\mu+1/2}^{(2)}(ka)} H_{\mu+1/2}^{(2)}(kr) .$$

It is known that

$$\text{Wronskian} \left\{ H_{\nu}^{(2)}(kr), H_{\nu}^{(1)}(kr) \right\} = - \frac{4i}{\pi kr} .$$



Hence:

$$C = \frac{1}{8A_\nu} \frac{H_{\nu+1/2}^{(2)}(ka)}{H_{\nu+1/2}^{(1)}(ka)}.$$

We may now write the total field:

$$u(r, \theta) = \frac{1}{8} \sum_{\nu} \frac{P_{\nu}(-\cos \theta)}{A_{\nu}} \frac{H_{\nu+1/2}^{(2)}(ka)}{H_{\nu+1/2}^{(1)}(ka)} \frac{H_{\nu+1/2}^{(1)}(kr >)}{\sqrt{r >}} \frac{Z_{\nu+1/2}(kr <)}{\sqrt{r <}}.$$

The sum is taken over all  $\nu$  such that  $P_{\nu}(-\cos \theta_0) = 0$ , and  $Z_{\nu+1/2}$  is defined by (3).

We now consider the diffraction of a plane wave by our obstacle. This is to be obtained by removing the point source to infinity in a suitable manner. Using the symbols in figure 2 (with  $\rho' = r'$ ), and letting the source go to  $-\infty$  along the x axis:

$$\frac{e^{ikR}}{4\pi R} \sim \frac{e^{ikr'}}{4\pi r'} e^{ikx}.$$

The total field  $v(r, \theta)$  from a plane wave  $e^{ikx}$  will be:

$$\begin{aligned} & \lim_{r' \rightarrow \infty} \frac{4\pi r'}{e^{ikr'}} u(r, \theta) \\ &= \lim_{r' \rightarrow \infty} \frac{4\pi r'}{e^{ikr'}} \sum_{\nu} \frac{P_{\nu}(-\cos \theta)}{A_{\nu}} \frac{H_{\nu+1/2}^{(2)}(ka)}{H_{\nu+1/2}^{(1)}(ka)} \frac{Z_{\nu+1/2}(kr)}{\sqrt{r}} \frac{H_{\nu+1/2}^{(1)}(kr')}{\sqrt{r'}}. \end{aligned}$$

Asymptotically

$$H_{\nu+1/2}^{(1)}(kr') \sim \sqrt{\frac{2}{\pi kr'}} e^{i(kr' - \frac{\nu\pi}{2} - \frac{\pi}{2})}.$$

The total field is:

$$(12) \quad v(r, \theta) = \sum_{\nu} \frac{P_{\nu}(-\cos \theta)}{A_{\nu}} \frac{H_{\nu+1/2}^{(2)}(ka)}{H_{\nu+1/2}^{(1)}(ka)} \frac{Z_{\nu+1/2}(kr)}{\sqrt{r}} \cdot \sqrt{\frac{\pi}{2k}} e^{-\frac{i\nu\pi}{2}}.$$

$A_{\nu}$  may be obtained from (see Sollfrey (5)):

$$\begin{aligned} A_{\nu} &= \int_{\theta_0}^{\pi} P_{\nu}^2(-\cos \theta) \sin \theta \, d\theta \\ &= -\frac{1}{2\nu+1} \sin^2 \theta_0 \left\{ \frac{\partial}{\partial(-\cos \theta)} P_{\nu}(-\cos \theta) \right\}_{\theta=\theta_0} \left\{ \frac{\partial}{\partial \nu} P_{\nu}(-\cos \theta) \right\}_{\theta=\theta_0}. \end{aligned}$$

In the case  $\theta_0 = \frac{\pi}{2}$ ,  $\nu$  is determined from  $P_{\nu}(0) = 0$ . Hence  $\nu$  is an arbitrary odd integer,  $2n+1$

$$A_{2n+1} = \int_{\frac{\pi}{2}}^{\pi} P_{2n+1}^2(-\cos \theta) \sin \theta \, d\theta = \int_0^1 P_{2n+1}^2(x) \, dx = \frac{1}{4n+3}.$$

We have

$$v(r, \theta) = \sum_{n=0}^{\infty} (4n+3) P_{2n+1}(-\cos \theta) \frac{H_{2n+3/2}^{(2)}(ka)}{H_{2n+3/2}^{(1)}(ka)} \frac{Z_{2n+3/2}(kr)}{\sqrt{r}} \sqrt{\frac{\pi}{2k}} (-1)^n.$$

This was obtained by Twersky (7) using a combination of an image method and superposition.

REFERENCES

- (1) W. Magnus, "Über die Beugung elektromagnetischer Wellen an einer Halbebene", Zeitschrift für Physik, 117, 1941, pp. 168-179.
- (2) A. Sommerfeld, Math. Annalen, 47, 1896, pp. 317-374.
- (3) D. S. Jones, "Removal of an Inconsistency in the Theory of Diffraction", Proceedings of the Cambridge Philosophical Society, 48, 1952, pp. 733-741.
- (4) M. J. Kontorowich and N. N. Lebedev, "On a Method of Solution of Some Problems of the Diffraction Theory", Journal of Physics, U.S.S.R., vol. I, No. 3, 1939, pp. 229-241.
- (5) W. Sollfrey, "Diffraction of Pulses by Conducting Wedges and Cones", N. Y. U., report EM-45, Math. Res Group.
- (6) S. Karp, "Diffraction by a Tipped Wedge", N. Y. U., report EM-52, Math. Res Group.
- (7) V. Twersky, J. Acoust. Soc. Am., 22, 539 (1950).

# STANFORD UNIVERSITY

## Technical Reports      Distribution List Contract Nonr 225(11) (NR-041-086)

Chief of Naval Research Office of Naval Research Department of the Navy Washington 25, D. C. Attn: Code 432	5	Office of Naval Research Department of the Navy Washington 25, D. C. Attn: Code 438	2
Scientific Section Office of Naval Research Department of the Navy 1000 Geary Street San Francisco 9, California Attn: Dr. J. D. Wilkes	2 1	Chief, Bureau of Ordnance Department of the Navy Washington 25, D. C. Attn: Re3d Re6a	1 1
Director, Naval Research Lab. Washington 25, D. C. Attn: Technical Information Officer	9	Chief, Bureau of Aeronautics Department of the Navy Washington 25, D. C.	1
Dr. Eugene Lukacs Head, Statistics Branch Office of Naval Research Department of the Navy Washington 25, D. C. Attn: Code 433	1	Chief, Bureau of Ships Asst. Chief for Research & Development Department of the Navy Washington 25, D. C.	1
Planning Research Division Deputy Chief of Staff Comptroller, U. S. A. F. The Pentagon Washington 25, D. C.	1	Director, ONR Branch Office 495 Summer Street Boston 10, Mass.	1
Headquarters, U. S. A. F. Director of Research and Development Washington 25, D. C.	1	Director, ONR Branch Office 844 N. Rush Street Chicago 11, Illinois	1
Chairman Research and Development Board The Pentagon Washington 25, D. C.	1	Director, ONR Branch Office 346 Broadway New York 13, N. Y.	1
Asst. Chief of Staff, G-4 for Research and Development U. S. Army Washington 25, D. C.	1	Director, ONR Branch Office 1030 E. Green St. Pasadena 1, California	1
Chief of Naval Operations Operations Evaluation Group- OP342E The Pentagon Washington 25, D. C.	1	Office of the Asst. Naval Attache' for Research Naval Attache' American Embassy Navy No. 100 Fleet Post Office New York, N. Y.	10

Commander, USNOTS  
Pasadena Annex  
3202 E. Foothill Boulevard  
Pasadena 8, California  
Attn: Technical Library 1

Office of Ordnance Research  
Duke University  
2127 Myrtle Drive  
Durham, North Carolina 1

Commanding Officer  
Ballistic Research Laboratory  
Aberdeen Proving Grounds  
Aberdeen, Maryland  
Attn: Mr. R. H. Kent 1

Director, David Taylor Model Basin  
Washington 25, D. C.  
Attn: L. Landweber 1  
P. Eisenberg 1  
Hydromechanics Lab. 1  
Technical Library 1

Director, National Bureau of  
Standards  
Washington, D. C.  
Attn: Natl. Hydraulics Lab. 1

Commanding General  
U. S. Proving Grounds  
Aberdeen, Maryland 1

Commander  
U. S. Naval Ordnance Test Station  
Inyokern, China Lake, Calif. 1

N. A. C. A.  
1724 F St., N.W.  
Washington 25, D. C.  
Attn: Chief, Office of Aero-  
nautical Intelligence 1

Hydrodynamics Laboratory  
National Research Laboratory  
Ottawa, Canada 1

Director  
Penn. State School of Engineering  
Ordnance Research Laboratory  
State College, Pa. 1

Statistical Laboratory  
Dept. of Mathematics  
University of California  
Berkeley 4, California 1

Hydrodynamics Laboratory  
California Inst. of Technology  
1201 East California St.  
Pasadena 4, California  
Attn: Executive Committee 1

Commanding Officer  
Naval Ordnance Laboratory  
White Oak, Silver Spring 19, Md.  
Attn: Technical Library 1

Ames Aeronautical Laboratory  
Moffett Field  
Mountain View, California  
Attn: Technical Librarian 1

Mr. Samuel I. Plotnick  
Asst. to the Director of Research  
The George Washington University  
Research Laboratory  
Area B, Camp Detrick  
Frederick, Maryland 1

Mathematics Library  
Syracuse University  
Syracuse 10, N. Y. 1

University of Southern California  
University Library  
3518 University Avenue  
Los Angeles 7, California 1

Library  
California Institute of Technology  
Pasadena 4, California 1

Engineering Societies Library  
29 W. 39th Street  
New York, N. Y. 1

John Crerar Library  
Chicago 1, Illinois 1

National Bureau of Standards  
Library  
3rd Floor North West Building  
Washington 25, D. C. 1

Library Mass. Inst. of Technology Cambridge 39, Mass.	1	Dr. E. P. Cooper U. S. Naval Shipyard U. S. Navy Radiological Defense Laboratory San Francisco, California	1
Louisiana State University Library University Station Baton Rouge 3, La.	1	Prof. R. Courant Inst. for Mathematics & Mechanics New York University New York 3, N. Y.	1
Library Fisk University Nashville, Tennessee	1	Dr. A. Craya Dept. of Aeronautical Engineering Columbia University New York 27, N. Y.	1
Mrs. J. Henley Crosland Director of Libraries Georgia Institute of Technology Atlanta, Georgia	1	Dr. K. S. M. Davidson Experimental Towing Tank Stevens Inst. of Technology 711 Hudson Street Hoboken, N. J.	1
Technical Report Collection Rm. 303A, Pierce Hall Harvard University Cambridge 38, Mass.	1	Prof. R. J. Duffin Dept. of Mathematics Carnegie Inst. of Technology Pittsburgh 13, Pa.	1
Prof. L. V. Ahlfors Dept. of Mathematics Harvard University Cambridge 38, Mass.	1	Dr. Carl Eckart Scripps Inst. of Oceanography La Jolla, California	1
Prof. Peter G. Bergmann Dept. of Physics Syracuse University Syracuse 10, N. Y.	1	Prof. A. Erdélyi Dept. of Mathematics California Institute of Technology Pasadena 4, California	1
Prof. G. Birkhoff Dept. of Mathematics Harvard University Cambridge 38, Mass.	1	Prof. K. O. Friedrichs Inst. for Mathematics & Mechanics New York University New York 3, N. Y.	1
Prof. H. Busemann Dept. of Mathematics University of So. California Los Angeles 7, California	1	Prof. D. Gilbarg Grad. Inst. for Applied Mathematics Indiana University Bloomington, Indiana	1
Dr. F. H. Clauser Dept. of Aeronautical Engineering Johns Hopkins University Baltimore 18, Maryland	1	Prof. A. T. Ippen Dept. of Civil and Sanitary Engineering Mass. Inst. of Technology Cambridge 39, Mass.	1
Dr. Milton Clauser Dept. of Aeronautical Engineering Purdue University Lafayette, Indiana	1		

Prof. Samuel Karlin Dept. of Mathematics Calif. Inst. of Technology Pasadena 4, California	1	Prof. C. B. Morrey Dept. of Mathematics University of California Berkeley 4, California	1
Prof. J. R. Kline Dept. of Mathematics University of Pennsylvania Philadelphia 4, Pa.	1	Prof. Z. Nehari Dept. of Mathematics Washington University St. Louis, Mo.	1
Dr. R. T. Knapp Hydrodynamics Laboratory California Inst. of Technology Pasadena 4, California	1	Dr. M. S. Plesset California Inst. of Technology Pasadena 4, California	1
Prof. P. A. Lagerstrom Aeronautics Dept. California Inst. of Technology Pasadena 4, California	1	Prof. W. Prager Dept. of Mathematics Brown University Providence 12, R. I.	1
Prof. Benjamin Lepson Mathematics Department Catholic University of America Washington 17, D. C.	1	Prof. P. C. Rosenbloom Dept. of Mathematics University of Minnesota Minneapolis 14, Minn.	1
Dr. Martin Lessen Aeronautical Engineering Dept. Penn. State College State College, Pa.	1	Dr. H. Rouse State Inst. of Hydraulic Research University of Iowa Iowa City, Iowa	1
Prof. H. G. Lew Dept. of Aeronautical Engineering Penn. State College State College, Pa.	1	Prof. Charles Saltzer Case Institute of Technology Cleveland, Ohio	1
Prof. Hans Lewy Dept. of Mathematics University of California Berkeley, California	1	Prof. A. C. Schaeffer Dept. of Mathematics University of Wisconsin Madison 6, Wisconsin	1
Prof. C. C. Lin Dept. of Mathematics Mass. Inst. of Technology Cambridge 39, Mass.	1	Prof. L. I. Schiff Dept. of Physics Stanford University Stanford, California	1
Prof. W. T. Martin Dept. of Mathematics Mass. Inst. of Technology Cambridge 39, Mass.	1	Prof. W. Sears Grad. School of Aeronautical Engineering Cornell University Ithaca, N. Y.	1
Dean Paul E. Mohn School of Engineering The University of Buffalo Buffalo, N. Y.	1	Prof. Herbert Solomon Teachers College Columbia University New York 27, N. Y.	1

Prof. D. Spencer Fine Hall, Box 708 Princeton, N. J.	1	Los Angeles Engineering Field Office Air Research and Development Command 5504 Hollywood Boulevard Los Angeles 28, California Attn: Capt. N. E. Nelson	1
Prof. J. J. Stoker Inst. for Mathematics & Mechanics New York University New York 3, N. Y.	1	Navy Department Naval Ordnance Test Station Underwater Ordnance Dept. Pasadena, California Attn: Dr. G. V. Schliestett Code P8001	1
Dr. V. L. Streeter Fundamental Mechanics Research Illinois Inst. of Technology Chicago 16, Illinois	1	ASTIA, Western Regional Office 5504 Hollywood Blvd. Hollywood 28, California	1
Prof. C. Truesdell Grad. Inst. for Applied Mathematics Indiana University Bloomington, Indiana	1	Armed Services Technical Information Agency Documents Service Center Knott Building Dayton 2, Ohio	5
Prof. Joseph Ullman Dept. of Mathematics University of Michigan Ann Arbor, Michigan	1	Mr. R. T. Jones Ames Aeronautical Laboratory Moffett Field Mountain View, California	1
Prof. J. K. Vennard Dept. of Civil Engineering Stanford University Stanford, California	1	Mr. J. D. Pierson Glenn L. Martin Company Middle River Baltimore, Maryland	1
Prof. Martin Vitousek University of Hawaii Honolulu 11, Hawaii	1	Mr. E. G. Straut Consolidated-Vultee Aircraft Corporation Hydrodynamics Research Lab. San Diego, California	1
Prof. S. E. Warschawski Dept. of Mathematics University of Minnesota Minneapolis 14, Minn.	1	Dr. George E. Forsythe National Bureau of Standards Institute for Numerical Analysis University of California 405 Hilgard Avenue Los Angeles 24, California	1
Prof. A. Weinstein Inst. for Fluid Dynamics & Applied Mathematics University of Maryland College Park, Maryland	1	Additional copies for project leader and assistants and re- serve for future requirements	50
Prof. A. Zygmund Dept. of Mathematics The University of Chicago Chicago 37, Illinois	1		
Mathematics Dept. University of Colorado Boulder, Colorado	1		